

There's Space Between Molecules

Mixing Liquids—Volume Is Not Conserved



Introduction

Mass and energy are conserved, but volume is not. Mixing alcohol and water demonstrates the principle—the particle model explain it.

Concepts

- Intermolecular forces—hydrogen bonding
- Polar molecules

Materials

Copper shot, Cu, 3–14 mesh, approximately 100 g	Graduated cylinders, 100-mL, 2
Ethyl alcohol, anhydrous, C_2H_5OH , 50 mL	Marbles, approximately 40–60
Sand, fine, approximately 100 g	Ping pong balls, approximately 10–15
Water, distilled or deionized	Stirring rod
Beaker, 600-mL	

Safety Precautions

Ethyl alcohol is a dangerous fire risk; it is flammable. The addition of denaturant (methyl alcohol) makes ethyl alcohol poisonous. Do not ingest. Clean up spills immediately. Wear chemical splash goggles and chemical-resistant gloves. Please review current Material Safety Data Sheets for additional safety, handling, and disposal information.

Procedure

Part A.

1. Fill a large beaker with ping pong balls. Ask students if the beaker is full.
2. Pour marbles into the beaker and repeat the question.
3. Repeat with the copper shot, the sand, and finally, the water.

Part B.

4. Carefully measure out exactly 50 mL of water in a 100-mL graduated cylinder.
5. Carefully measure out exactly 50 mL of anhydrous ethyl alcohol into a second 100-mL graduated cylinder.
6. Pour the water from the first graduated cylinder into the graduated cylinder containing the ethyl alcohol.
7. Stir the mixture of alcohol and water with a stirring rod and wait about one minute for the bubbles to come out of solution.
8. Observe that the final volume of liquid in the cylinder is less than 100 mL.

Disposal

Please consult your current *Flinn Scientific Catalog/Reference Manual* for general guidelines and specific procedures governing the disposal of laboratory waste. The resulting solution may be flushed down the drain with excess water according to Flinn Suggested Disposal Method #26b.

Tips


- Part B of this demonstration is even more spectacular if done in a 24" glass demonstration tube (Flinn Catalog No. GP9146). Fill the tube with equal volumes of deionized water (optionally colored with food dye) and anhydrous ethyl alcohol, stopper the ends and begin to mix the solvents by turning the tube. An air bubble will soon appear out of nowhere. Adding equal volumes of water and ethyl alcohol to a volumetric flask also works well.
- Anhydrous ethyl alcohol does not contain any water. Do not use 95% alcohol that contains 5% water.

Discussion

A beaker filled to the brim with objects such as ping pong balls or marbles may seem completely packed. But this demonstration shows there is still space between the objects that can be filled with progressively smaller objects. Even liquids such as ethyl alcohol and water have space between the molecules.

When 50 mL of water is added to 50 mL of water or when 50 mL of alcohol is added to 50 mL of alcohol, the final volume will always be 100 mL, as expected. In this demonstration, when the water is added to the alcohol, the final volume is about 10% less than the original volume of the two liquids. The "vanishing volume" is due to differences in packing of the solvent molecules in the mixture versus the pure substances. Molecules of ethyl alcohol actually pack together more closely with water molecules than with other alcohol molecules due to hydrogen bonding. The solvent molecules form a highly-laced, 3-dimensional network held together by strong hydrogen bonds (Figure 1). Each alcohol molecule is able to form as many as three hydrogen bonds with neighboring water or alcohol molecules. The result is an intricate lattice or network of molecules strongly attracted to one another.

Looking at the Molecules



Hydrogen bonding is an especially strong form of dipole-dipole interaction. A dipole-dipole interaction is the attraction of the positive end of one polar molecule for the negative end of another polar molecule. In hydrogen bonding, a hydrogen atom serves as a bridge between two electronegative atoms (nitrogen, oxygen, or fluorine).

Hydrogen bonding plays a major role in the properties of water and alcohols. Hydrogen bonding between water molecules leads to a very high boiling point when compared to other similar liquids. The effect of hydrogen bonding can also be clearly seen when boiling points for alcohols are compared to nonpolar ethers having the same molecular weight. Consider butyl alcohol and ethyl ether. Both have the same formula ($C_4H_{10}O$), the same molecular weight (74 g/mole), and the same size. Butyl alcohol, however, boils at 118 °C, while diethyl ether boils at 35 °C. The 80°C difference in boiling points is due to the hydrogen bonding in

Figure 1. Hydrogen bonding between alcohol and water.

the butyl alcohol.

Connecting to the National Standards

This laboratory activity relates to the following National Science Education Standards (1996):

Unifying Concepts and Processes: Grades K–12

Evidence, models, and explanation
Constancy, change, and measurement

Content Standards: Grades 5–8

Content Standard B: Physical Science, properties and changes of properties in matter

Content Standards: Grades 9–12

Content Standard B: Physical Science, structure of atoms, structure and properties of matter, chemical reactions

Flinn Scientific—Teaching Chemistry™ eLearning Video Series

A video of the *There's Space Between Molecules* activity, presented by Jamie Benigna, is available in *Mixing Liquids—Volume Is Not Conserved*, part of the Flinn Scientific—Teaching Chemistry eLearning Video Series.

Materials for *There's Space Between Molecules* are available from Flinn Scientific, Inc.

Catalog No.	Description
C0083	Copper Shot, 3–14 mesh, 100 g
S0003	Sand, Fine, White, 500 g
E0012	Ethyl alcohol, anhydrous, 500 mL
GP2020	Graduated cylinder, glass, 100-mL
GP9146	Demonstration tube, glass, 249

Consult your *Flinn Scientific Catalog/Reference Manual* for current prices.